



Carbohydrates I, II and III

Dr. Diala Abu-Hassan, DDS, PhD

Dr.abuhassand@gmail.com

Medical students- summer semester
2015

All images were taken from Campbell
textbook except where noted

What Are Carbohydrates?

Carbohydrates are members of a large class of naturally occurring polyhydroxy ketones and aldehydes

Roles of Carbohydrates



Major energy sources

Oligosaccharides play a key role in **cell–cell interactions** and **immune recognition**.

Polysaccharides are essential **structural components** of several classes of organisms such as cellulose (a major component of grass and trees)

Aldoses and ketoses


- 1 A comparison of glyceraldehyde (an aldotriose) and dihydroxyacetone (a ketotriose).




Nomenclature: aldo- or keto- + latin number + ose

Most common sugars are aldoses rather than ketoses

Optical isomers (stereoisomers)



D-Glyceraldehyde




L-Glyceraldehyde

Enantiomers

More Cs, more stereoisomers

D sugars, rather than L sugars, predominate in nature


How to determine chiral carbons?



Stereoisomers of an aldotetrose

Number of stereoisomers = 2^n , n is the number of chiral carbons

A Diastereomers
D-erythrose and
D-threose.



Stereoisomers of an aldotetrose

Diastereomers: non-superimposable, non-mirror image stereoisomers





FIGURE 16.3 Stereoisomers of an aldotetrose.

Diastereomers


Epimers are diastereomers that differ in the configuration at only **ONE** chiral carbon
D-erythrose and D-threose are epimers.

Stereoisomers


A Aldoses containing from three to six carbon atoms, with the numbering of the carbon atom shown. Note that the figure shows only half the possible isomers. For each isomer shown, there is an enantiomer that is not shown, the L series.




Structure of Glucose



Structure of Glucose




Glucose anomers



Fischer projection formulas of three forms of glucose.




Glucose anomers and cyclization




In some molecules, any anomer can be used

In other sugars, only one anomer is used, e.g. only β -D-ribose and β -D-deoxyribose are found in RNA and DNA, respectively.

Structural representation of sugars



Structural Representation of Sugars




Carbohydrates

- **Mono- (one), Oligo- (a few) and (many) Polysaccharides**
- **Monosaccharides (simple sugar)** a carbohydrate with three to seven carbon atoms. General formula $C_n(H_2O)_n$.
- A monosaccharide can be a polyhydroxy aldehyde (**aldose**) or a polyhydroxy ketone (**ketose**)
- **Aldose A monosaccharide that** contains an aldehyde carbonyl group.
- **Ketose A monosaccharide that** contains a ketone carbonyl group
- Have many hydroxyl groups on adjacent carbons together with either an aldehyde or ketone group

Glucose

- The most important simple carbohydrate in human metabolism.
- It is the final product of carbohydrate digestion and provides acetyl groups for entry into the citric acid cycle as acetyl-SCoA

Structure of Glucose




Galactose

- Plant gums and pectins
- A component of the disaccharide lactose (milk sugar)
- Is produced from lactose during digestion.
- Is converted to glucose to provide energy
- Is synthesized from glucose to produce lactose for milk and compounds needed in brain tissue.



Jam with galactose in the pectin that stiffens it

Galactose




An aldohexose


It differs from glucose only in the orientation of the OH group at C4.

Galactose vs Glucose

Galactose




Glucose



Fructose



- D-Fructose, often called levulose or fruit sugar
- honey and many fruits.
- one of the two monosaccharides combined in the disaccharide sucrose.
- Fructose is produced commercially in large quantities by hydrolysis of cornstarch to make high fructose corn syrup (HFCS).

Fructose



- Like glucose and galactose, fructose is a 6-carbon sugar.
- A ketohexose
- It forms a five-membered rings in solution

Ribose and 2-Deoxyribose




- Both are 5-carbon aldehyde sugars.
- Important as parts of larger biomolecules, such as coenzyme A , ATP, cyclic AMP and in oxidizing and reducing agent coenzymes.
- 2-deoxyribose differs from ribose by the absence of one oxygen atom, that in the OH group at C2.
- Both exist as mixtures of open-chain and cyclic hemiacetal forms.

Sugar derivatives

Amino sugars

An amino group ($-\text{NH}_2$) or one of its derivatives is substituted for the hydroxyl group of the parent sugar.



Components of bacterial cell walls


Reactions of Monosaccharides

Oxidation-Reduction Reactions

Oxidation

Aldehyde		Carboxylic acid
Cyclic aldose		Lactone
Ketose		Aldose

Oxidation of a sugar to a lactone




- Carbohydrates that react with oxidizing agents are classified as **reducing sugars** (they reduce the oxidizing agent).
- Glucose** can be detected **specifically** by the enzyme **glucose oxidase**.



■ **FIGURE 16.10** A silver mirror produced by an aldehyde. After the addition of Tollens reagent to an aldehyde, a silver mirror has been deposited in the inside of this flask.


Ketoses can become aldoses

- In basic solution, ketoses are also reducing sugars.
- Rearrangement.




Reduced sugars (deoxy sugars)

L-fucose (L-6-deoxygalactose) is found in the carbohydrate portions of some glycoproteins including the ABO blood-group antigens.




β -L-Fucose
(6-Deoxy- β -L-galactose)




β -L-Galactose

D-2-deoxyribose is found in DNA




β -D-Deoxyribose
(2-Deoxy- β -D-ribose)




β -D-Ribose


Reduced sugars (polyhydroxy alcohols or alditols)




D-Sorbose



D-Sorbitol



D-Xylitol




D-Xylulose

Sorbitol and xylitol are used commercially as sweeteners in sugarless chewing gum and candy.

Reactions of Monosaccharides

Esterification of sugars




Phosphate esters formation (phosphorylated sugars)

Important for the metabolism of sugars

Reactions of Monosaccharides

Reaction with Alcohols: Glycoside Formation


Hemiacetals react with alcohols with the loss of water to yield acetals



- Monosaccharides are cyclic hemiacetals that react with alcohols to form acetals (**glycosides**).
- O-glycosides not N-glycosides
- **Furanosides** are glycosides derived from furanoses, and **pyranosides** are glycosides derived from pyranoses.

Hydrolysis of a disaccharide


- Hydrolysis reaction takes place during digestion of all carbohydrates.



Reactions of Monosaccharides

Glycosidic bond between monosaccharides


2 glucose molecules



Chemical natures of oligo-and polysaccharides depend on:

Monosaccharides linked together


The type of glycosidic bond formed (anomer type and carbons linked)



Linear and branched-chain polymers of α -D-glucose


A

The linear polyglucose chain occurs in amylose. All glycosidic bonds are $\alpha(1\rightarrow4)$.




B

The branched-chain polymer occurs in amylopectin and glycogen. Branched-polyglucose-chain glycosidic bonds are $\alpha(1\rightarrow6)$ at branch points, but all glycosidic bonds along the chain are $\alpha(1\rightarrow4)$.



Polysaccharides and reducing sugars




■ FIGURE 16.17 Reducing sugars.

A disaccharide with a free hemiacetal end is a reducing sugar because of the presence of a free anomeric aldehyde carbonyl or potential aldehyde group.


Oligosaccharides

Disaccharides are Oligosaccharides



Sucrose

- **Common table sugar, cane sugar, beet sugar**
- Hydrolysis of sucrose yields one molecule of D-glucose and one molecule of D-fructose.
- The 50:50 mixture of glucose and fructose (invert sugar) is sweeter than sucrose and is used as a food additive.
- No hemiacetal group because a 1,2 link joins both anomeric carbon atoms.
- Sucrose is not reducing because it does not have a hemiacetal group
- Sucrose is the only common disaccharide that is not a reducing sugar.



Sucrose (glucose- α -1,2-fructose)

Biochemical Application: Synthetic sugars and sweetness

TABLE 22.1 Relative Sweetness of Some Sugars and Sugar Substitutes


NAME	TYPE	SWEETNESS
Lactose	Disaccharide	16
Galactose	Monosaccharide	30
Maltose	Disaccharide	33
Glucose	Monosaccharide	75
Sucrose	Disaccharide	100
Fructose	Monosaccharide	175
Cyclamate	Artificial	3000
Aspartame	Artificial	15,000
Saccharin	Artificial	35,000
Sucralose	Artificial	60,000

Biochemical application: Sucralose

Sucralose is not metabolized by the body (does not provide calories).

So far, it is a safe sugar substitute.

Modified galactose and fructose




Sucralose

■ **FIGURE 16.20** The structure of sucralose. Note that sucralose (marketed under the trade name Splenda) differs from sucrose in the substitution of chlorine for three hydroxyls.

Lactose (milk sugar)

- Human milk is about 7% lactose.
- A disaccharide of β -D-galactose and either α or β -D-glucose.
- The two monosaccharides are connected by a β -1,4 link.
- A reducing sugar because the glucose ring (on the right) is a hemiacetal at C1.



Clinical hint-Lactose intolerance



© Thomson Learning/Charles D. Winters

- Is prevalent in adults in all populations.
- Due to lactase enzyme deficiency.
- Because lactose remains in the intestines rather than being absorbed, it raises the osmolarity, which draws in excess water.
- Intestinal bacteria ferment lactose to produce lactate, carbon dioxide, hydrogen gas, and methane resulting in bloating, cramps, and diarrhea.
- Lactose free food, the use of commercial enzyme preparations and Lactaid, milk that has been treated with lactase to reduce its lactose content


■ **Dairy substitutes for the lactose intolerant.**
These products help those with lactose intolerance meet their calcium needs.

Maltose (Malt sugar)

- Present in fermenting grains
- Can be prepared by enzyme-catalyzed degradation of starch.
- Used in prepared foods as a sweetener.
- Produced during starch digestion by α -amylase in the small intestine
- Hydrolyzed to glucose by maltase.

Maltose


- Two -D-glucose molecules are joined in maltose by an α -1,4 link
- It is both an acetal (at C1 in the left-side glucose) and a hemiacetal (at C1 in the right-side glucose).
- The hemiacetal group on the right can react with an oxidizing agent, thus maltose can be a reducing sugar.



Maltose (glucose- α -1,4-glucose)

Cellobiose

- A disaccharide that is obtained from the hydrolysis of cellulose
- Two -D-glucose molecules are joined by a **β -1,4 link**
- It is both an acetal (at C1 in the left-side glucose) and a hemiacetal (at C1 in the right-side glucose).
- The hemiacetal group on the right can react with an oxidizing agent, thus maltose can be a reducing sugar.




Cellobiose (glucose- β -1,4-glucose)

Polysaccharides


- Homopolysaccharide: a polymer that consists of only one type of monosaccharide
- Heteropolysaccharide a polymer that consists of more than one type of monosaccharide
- Glucose is the most common monomer

Cellulose





- Animals lack cellulases enzymes that hydrolyze cellulose to glucose.
- Cellulases are found in the bacteria that inhabit the digestive tracts of insects and grazing animals, such as cattle and horses.

H-bonds in cellulose




Starch



Starches are polymers of α -D-glucose that occur in plant cells, usually as starch granules in the cytosol.


Types of starches

Amylose



α (1-4)

Types of starches




Amylose forms a helix with six residues per turn.

Iodine molecules can fit inside the helix to form a starch–iodine complex (dark-blue color)

Types of starches

Amylopectin

No known preferred conformation for amylopectin
Branch points occur about every 25 residues



Amylopectin and glycogen react with iodine to produce a red-brown product, not blue.

Amylases

- Starches are storage molecules
- Both plants and animals contain enzymes that hydrolyze starches (α - and β -amylase)
- Both enzymes attach $\alpha(1-4)$ linkages


α - amylase	β -amylase
Endoglycosidase	Exoglycosidase
Acts on glycosidic linkages along the chain	Acts on the nonreducing end of the polymer
Produces glucose and maltose	Produces maltose (glucose dimer)

Amylase action on amylose and amylopectin

- Amylose can be completely degraded to glucose and maltose by the two amylases
- Amylopectin is not completely degraded by the two amylases because the branching linkages are not attacked.
- Debranching enzymes degrade the $\alpha(1-6)$ linkages

Glycogen




- glycogen is more highly branched than starch
- Branch points occur about every 10 residues in glycogen
- α (1-4) linkage in main chain
- α (1-6) linkage at branches
- The average chain length is 13 glucose residues
- 12 layers of branching



Glycogen


Glycogen

- Glycogen is found in granules in animal cells particularly in well-fed liver and muscle cells
- Some athletes (long-distance runners) try to build up their glycogen reserves before a race by eating large amounts of carbohydrates.
- Degradative enzymes remove glucose units from glycogen when energy is needed
- Glycogen phosphorylase cleaves one glucose at a time from the non-reducing end of a branch to produce glucose-1-phosphate




Glycogen


- Debranching enzymes complete the breakdown of glycogen.
- The number of branch points is important because:
 1. A more branched polysaccharide is more water soluble.
 2. When energy is needed quickly, the glycogen phosphorylase has more potential targets if there are more branches, allowing a quicker mobilization of glucose.



Amylopectin versus glycogen



Amylopectin




Glycogen

- Branching
- Source



Pectin

- A polysaccharide of plant cell wall
- A polymer made up mostly of D-galacturonic acid
- D-galacturonic acid is a galactose derivative
- A gelling agent in yogurt, fruit preserves, jams, and jellies.



D-Galacturonic acid

Chitin



Has a structural role


Mechanical strength because of H-bonds between strands.

A major structural component of the exoskeletons of invertebrates such as insects and crustaceans (a group that includes lobsters and shrimp), and the cell walls of algae, fungi, and yeasts.

Polysaccharides in bacterial cell wall

- Heteropolysaccharides are major components of bacterial cell walls


A



- N*-Acetylmuramic acid is found only in prokaryotic cell walls; it does not occur in eukaryotic cell walls.

Polysaccharides in bacterial cell wall

Peptidoglycan is the material that results from the crosslinking of polysaccharides by peptides



Fibers and health

- Stimulate peristaltic action
- Bind to toxic substances in foods they get exported from the body, thus preventing them from damaging the lower intestine or being reabsorbed there
- Bind carcinogens, thus prevent cancer
- Bind cholesterol resulting in less amount in the blood
- Fewer calories




Glycoseaminoglycans (GAGs)

Polysaccharides that have a repeating disaccharide

One of the sugars is an amino sugar


At least one of the sugars has a negative charge due to the presence of a sulfate group or a carboxyl group

GAGs are linked to proteins to form proteoglycans







Heparin is a natural anticoagulant

Hyaluronic acid is a component of the vitreous humor of the eye and of the lubricating fluid of joints




Glycoseaminoglycans (GAGs)

GAGs are highly polar and attract water, thus act as a lubricant or as a shock absorber




Glycoproteins

- Glycoproteins contain carbohydrate residues attached to the polypeptide chain
- Example: antibodies.



Carbohydrates as **antigenic determinants**

- Carbohydrates act as **antigenic determinants**, the portions of an antigenic molecule that antibodies recognize and bind.



■ **FIGURE 16.30** The structures of the blood-group antigenic determinants.

Glycoprotein and blood transfusion

Transfusion Relationships			
Blood Type	Makes Antibodies Against	Can Receive From	Can Donate To
O	A, B	O	O, A, B, AB
A	B	O, A	A, AB
B	A	O, B	B, AB
AB	None	O, A, B, AB	AB

Glycoprotein versus Proteoglycan

Low carbohydrate diet



- More carbohydrates, more insulin, more fat synthesis and storage and inhibition of fat burning