

PHYSIOLOGY

☐ Sheet

☒ Slide

☐ Handout

Number

11

Subject


The Eye: III. Central Neurophysiology of Vision

Doctor

Faisal Mohammed

Date: 00/00/2016

Price:



The Eye: III. Central Neurophysiology of Vision L12

Faisal I. Mohammed, MD, PhD



Objectives


- List the stations for the visual pathway
- Determine the functions of the visual cortices
- Describe visual neural processing
- Explain the abnormalities of visual pathway lesions
- Outline the importance of the light reflexes direct and consensual





Visual Pathways to the Brain

- ▶ optic nerve
 - ▶ axons of ganglion cells of the retina
- ▶ optic chiasm
 - ▶ all fibers from the nasal halves of the retina cross to the opposite side and join fibers from the opposite temporal retina to form the optic tracks
 - ▶ Fibers project to the pretectal area and superior colliculli
- ▶ synapse in the dorsal lateral geniculate nucleus (LGN) of the thalamus
- ▶ from LGN to primary visual cortex by way of the optic radiation

Visual pathways to the brain









Visual field of left eye

Visual field of right eye






C. Visual field loss		Name of deficit
left eye	right eye	
		0 Normal vision
		1 Anopsia of right eye (Blindness)
		2 Heteronymous (Bitemporal) hemianopsia
		3 Homonymous hemianopsia
		4 Upper quadrant anopsia
		5 Lower quadratic anopsia with macular sparing
		6 Upper quadratic anopsia with macular sparing

I

Neural pathway for vision






Retinal Projections to Subcortical Regions

- ▶ suprachiasmatic nucleus of the hypothalamus
 - ▶ control of circadian rhythms??
- ▶ pretectal nuclei
 - ▶ pupillary light reflex
 - ▶ accommodation of the lens
- ▶ superior colliculus
 - ▶ rapid directional movement of both eyes
- ▶ ventral lateral geniculate
 - ▶ control of bodies behavioral functions??

Autonomic Pathways to the Eye





Function of the Dorsal Lateral Geniculate

- ▶ Two principle functions:
 - ▶ relay of information to primary visual cortex
 - ▶ “gate control” of information to primary visual cortex



“Gate” Function of the LGN


- ▶ LGN receives input from corticofugal fibers originating in the primary visual cortex.
- ▶ Input from reticular areas of the midbrain
- ▶ Both inputs are inhibitory and can turn off transmission of the signal in select areas of the LGN.
- ▶ Both inhibitory inputs presumably control the visual input that is allowed to pass to the cortex.



Primary Visual Cortex

- ▶ located in the occipital lobe in the calcarine fissure region
- ▶ large representation in visual cortex for the macula (region for highest visual acuity)
- ▶ layered structure like other cortical areas
- ▶ columnar organization as well
- ▶ receives the primary visual input

Location of Visual Areas in the Cortex






Secondary Visual Areas

- ▶ visual association cortex
- ▶ responsible for analyzing the visual information
- ▶ area for 3 dimensional position, gross form, and motion
- ▶ area for color analysis



Processing in the Visual Cortex

- ▶ separation of the signals from the two eyes is lost in the primary visual cortex
- ▶ signals from one eye enter every other column, alternating with signals from the other eye
- ▶ allows the cortex to decipher whether the two signals match



Connections in the Visual Cortex



Analysis of the Visual Image

- ▶ The visual signal in the primary visual cortex is concerned mainly with contrasts in the visual scene.
- ▶ The greater the sharpness of the contrast, the greater the degree of stimulation.
- ▶ Also detects the direction of orientation of each line and border.
 - ▶ for each orientation of a line, a specific neuronal cell is stimulated.



Fields of Vision


- ▶ nasal field located on the temporal side of the retina
- ▶ temporal field located on the nasal side of the retina
- ▶ interruption of the visual pathway at different points leads to very specific visual field defects
- ▶ “blind spot” located about 15 degrees lateral to the central point of vision
- ▶ no rods or cones in this area, called the optic disc, exit point for axons of the ganglion cells




The Autonomic Nerves to the Eyes

- ▶ The eye is innervated by both parasympathetic and sympathetic neurons.
- ▶ Parasympathetic fibers arise in the Edinger-Westphal nucleus, pass in the 3rd cranial nerve to the ciliary ganglion.
 - ▶ Postganglionic fibers excite the ciliary muscle and sphincter of the iris.
- ▶ Sympathetic fibers originate in the intermediolateral horn cells of the superior cervical ganglion.
 - ▶ Postganglionic fibers spread along the carotid artery and eventually innervate the radial fibers of the iris.

Autonomic Pathways to the Eye



Pupillary reflex pathway (Direct and Consensual –Indirect)





Control of Accommodation (Focusing the Eyes)

- ▶ results from contraction or relaxation of the ciliary muscle
- ▶ regulated by negative feedback mechanism that automatically adjust the focal power of the lens for highest degree of visual acuity within about 1 sec
- ▶ exact mechanism is not known



Control of Pupillary Diameter

- ▶ *miosis*: decreasing of pupillary aperture due to stimulation of parasympathetic nerves that excite the pupillary sphincter muscle
- ▶ *mydriasis*: dilation of pupillary aperture due to stimulation of sympathetic nerves that excite the radial fibers of the iris



Pupillary Light Reflex

- ▶ When the amount of light entering the eyes increases, the pupils constrict.
- ▶ Functions to help the eye adapt extremely rapidly to changing light conditions.
- ▶ Light excites fibers going to pretectal nuclei.
- ▶ From pretectal nuclei fibers pass to Edinger-Westphal nucleus and back through parasympathetic nerves to constrict iris sphincter.

... |

Thank You

